arXiv:1508.03415v1 [cs.IT] 14 Aug 2015

SEVERAL CLASSES OF BENT, NEAR-BENT AND 2-PLATEAUED FUNCTIONS OVER FINITE FIELDS OF ODD CHARACTERISTIC

GUANGKUI XU, XIWANG CAO

Abstract. Inspired by a recent work of Mesnager, we present several new infinite families of quadratic ternary bent, near-bent and 2-plateaued functions from some known quadratic ternary bent functions. Meanwhile, the distribution of the Walsh spectrum of two class of 2-plateaued functions obtained in this paper is completely determined. Additionally, we construct the first class of p-ary bent functions of algebraic degree p over the fields of an arbitrary odd characteristic. The proposed class contains non-quadratic p-ary bent functions that are affinely inequivalent to known monomial and binomial ones.

1. Introduction

Boolean functions with few Walsh transform values have useful applications in cryptography and communications. Such functions provide protection against linear cryptanalysis in cryptography [24] and correspond to sequences that have low cross-correlation with the m-sequence in communications [14]. Boolean bent functions which have the highest possible nonlinearity with even number of variables were first introduced by Rothaus [29] in 1976. Bent functions have been widely studied and have received much attention due to their significantly important role in cryptography, coding theory, communication, and sequence design. As a generalization of Rothau's notion of a bent function, Kumar, Scholtz and Welch extended it to p-ary bent functions from $\mathbb{Z}_p^n$ to $\mathbb{Z}_p$ in [20], where p is an integer. In the case of p-ary bent functions things are naturally much more complicated compare to Boolean bent functions. A number of recent papers are devoted to the description of new classes of bent functions. However, there are a few known families of bent functions. In general, there are two ways to construct bent functions: primary constructions and secondary constructions. For some primary and secondary constructions of bent functions on monomials, binomials and quadratic functions, the reader is referred to [1, 2, 6, 12, 16, 21, 22, 25]. Surveys of known results on bent functions can be seen in [4, 13, 28] and the references therein. Plateaued functions were introduced by Zheng and Zhang as good candidates for designing cryptographic functions since they possess desirable various cryptographic characteristics [36]. For more results on the treatment of plateaued functions, we refer to [5, 7, 8, 9, 10, 19, 26, 35, 37]

For a prime p and a positive integer n, let $\mathbb{F}_{p^n}$ be the finite field with p^n elements and $\mathbb{F}_{p^n}^* = \mathbb{F}_{p^n} \setminus \{0\}$. For any k dividing n, we denote the trace function from $\mathbb{F}_{p^n}$ to $\mathbb{F}_{p^k}$ as follows:

$$\mathrm{Tr}_k^n(x) = x + x^{p^k} + x^{p^{2k}} + \cdots + x^{p^{n-k}}.$$

For $k = 1$, $\mathrm{Tr}_1^n(x) = \sum_{i=0}^{n-1} x^{p^i}$ is called the absolute trace function. Recently, Mesnager [27] proved a strong version of [3, Theorem 3], and provided several primary and secondary constructions of bent functions. Via some known binary monomial bent functions and their

---

1991 *Mathematics Subject Classification.* (MSC 2010) 94A60.

Key words and phrases. p-ary bent function; near-bent function; 2-plateaued function; Walsh transform.

corresponding dual functions, she obtained two new infinite families of bent functions with the forms

$$f(x) = \mathrm{Tr}_1^k(\lambda x^{2^k+1}) + \mathrm{Tr}_1^n(ux)\mathrm{Tr}_1^n(vx) \tag{1.1}$$

and

$$f(x) = \mathrm{Tr}_1^k(x^{2^k+1}) + \mathrm{Tr}_1^n\left(\sum_{i=1}^{2^{r-1}-1} x^{(2^k-1)\frac{i}{2^r}+1}\right) + \mathrm{Tr}_1^n(ux)\mathrm{Tr}_1^n(vx) \tag{1.2}$$

over $\mathbb{F}_{2^n}$, where $n = 2k$, $\lambda \in \mathbb{F}_{2^k}^*$ and $u, v \in \mathbb{F}_{2^n}^*$. It is well known that a function given by adding a linear function to one bent function is also bent. Inspired by the work of [27], a very natural question is considered: whether we can obtain new p-ary bent functions by adding the product of two linear functions to some known p-ary bent functions? In a very recent paper [33], the authors proved that some such p-ary functions are also bent under certain conditions. In this paper, we will continue the work of [27, 33] and construct more ternary bent and 2-plateaued functions of the form

$$f(x) = \mathrm{Tr}_1^k(\lambda x^{3^k+1}) + \mathrm{Tr}_1^n(ux)\mathrm{Tr}_1^n(vx) \tag{1.3}$$

over $\mathbb{F}_{3^n}$, where $n = 2k$, $\lambda \in \mathbb{F}_{3^k}^*$ and $u, v \in \mathbb{F}_{3^n}^*$. For an odd prime p, some p-ary bent functions of the form

$$f(x) = \mathrm{Tr}_1^n(\lambda x^d) - \mathrm{Tr}_1^n(ux)\mathrm{Tr}_1^n(x)^{p-1} \tag{1.4}$$

over $\mathbb{F}_{p^n}$ are obtained, where $d = p^k + 1$ or 2. The main result of this paper is obtained from the study of the Walsh transform, which is different from the one used in [27].

The rest of the paper is organized as follows. In Section 2, we give some notations and related results. In Section 3, using some known ternary bent functions, we derive more ternary bent, near-bent and 2-plateaued functions of the form (1.3). In Section 4, two new classes of non-quadratic p-ary bent functions of algebraic degree p with the form (1.4) are obtained.

2. PRELIMINARIES

A polynomial $f(x) \in \mathbb{F}_{p^n}[x]$ is called a *permutation polynomial* of $\mathbb{F}_{p^n}$ if it induces a bijective map from $\mathbb{F}_{p^n}$ to $\mathbb{F}_{p^n}$. For a permutation polynomial $f(x) \in \mathbb{F}_{p^n}[x]$ there exists (a unique) $f^{-1}(x) \in \mathbb{F}_{p^n}[x]$ such that $f(f^{-1}(x)) \equiv f^{-1}(f(x)) \equiv x \pmod{x^{p^n} - x}$. We call $f^{-1}(x)$ the *compositional inverse* of $f(x)$. A polynomial of the form

$$L(x) = \sum_{i=0}^{m} a_i x^{p^i} \in \mathbb{F}_{p^n}[x]$$

is called a *linearized polynomial*. It is well known that the compositional inverse of a linearized permutation polynomial is also a linearized polynomial. Recently, the compositional inverse of of some linearized permutation polynomials have been discovered. For detailed information, the reader is referred to [11, 30, 31, 32].

For some $l \mid n$, denote by $N_l^n : \mathbb{F}_{p^n} \to \mathbb{F}_{p^l}$ the norm function $N_l^n(x) = x^{\frac{p^n-1}{p^l-1}}$.

Lemma 1 (Theorem 2.1 [31]). *Let $a \in \mathbb{F}_{p^n}^*$ and $d = \gcd(n, r)$. Then the linearized binomial $L_{a,r}(x) = x^{p^r} + ax$ is a permutation polynomial over $\mathbb{F}_{p^n}$ if and only if $(-1)^{\frac{n}{d}} N_d^n(a) \neq 1$. Moreover,*

$$L_{a,r}^{-1}(x) = \frac{N_d^n(a)}{N_d^n(a) + (-1)^{\frac{n}{d}}} \sum_{i=0}^{\frac{n}{d}-1} (-1)^i a^{-\frac{p^{(i+1)r}-1}{p^r-1}} x^{p^{ir}}.$$

In particular, if n is even, then $L_{a,\frac{n}{2}}(x) = x^{p^{\frac{n}{2}}} + ax$ is a linearized permutation polynomial over $\mathbb{F}_{p^n}$ if and only if $a^{p^{\frac{n}{2}}+1} \neq 1$, and

$$L^{-1}_{a,\frac{n}{2}}(x) = \frac{1}{a^{p^{\frac{n}{2}}+1} - 1}(a^{p^{\frac{n}{2}}} x - x^{\frac{n}{2}}).$$

Let $f : \mathbb{F}_{p^n} \to \mathbb{F}_p$ be a p-ary function with n variables. For any integer $d \in \{0, 1, \cdots, p^n - 1\}$, let $\sum_{i=0}^{n-1} d_i p^i$ be its p-ary expansion with $0 \leq d_i \leq p-1$, then the *p-weight* of d denoted by $w_p(d)$ equals $\sum_{i=0}^{n-1} d_i$. Recall that each function $f : \mathbb{F}_{p^n} \to \mathbb{F}_p$ can be represented by a univariate polynomial over $\mathbb{F}_{p^n}$. The algebraic degree of $f(x)$ is equal to the maximum p-weight of an exponent j of the term $a_j x^j$ in $f(x)$ with $a_j \neq 0$.

The Walsh transform of f is defined as follows

$$\widehat{\chi}_f(a) = \sum_{x \in \mathbb{F}_{p^n}} \omega^{f(x) - \mathrm{Tr}_1^n(ax)}, a \in \mathbb{F}_{p^n},$$

where $\omega = e^{2\pi\sqrt{-1}/p}$ is a primitive p-th root of unity. The values $\widehat{\chi}_f(a), a \in \mathbb{F}_{p^n}$ are called the *Walsh coefficients* of f. The *Walsh spectrum* of a Boolean function f is the multiset $\{\widehat{\chi}_f(a), a \in \mathbb{F}_{p^n}\}$.

Definition 1. *A p-ary function f is called bent if $|\widehat{\chi}_f(a)| = p^{n/2}$ for all $a \in \mathbb{F}_{p^n}$.*

A p-ary bent function $f(x)$ is called *regular* if $\widehat{\chi}_f(a) = p^{n/2}\omega^{f^*(a)}$ for any $a \in \mathbb{F}_{p^n}$, where the function $f^*(x)$ is called the dual of $f(x)$. A bent function $f(x)$ is called *weakly regular* if there is a complex μ with unit magnitude such that $\widehat{\chi}_f(a) = p^{n/2}\mu\omega^{f^*(a)}$. The dual of a (weakly) regular bent function is again a (weakly) regular bent function [15].

Definition 2. [7] *For an integer $0 \leq s \leq n$, if $|\widehat{\chi}_f(a)| = \{0, p^{\frac{n+s}{2}}\}$ for all $a \in \mathbb{F}_{p^n}$, then we call f s-plateaued (for $s = 1$ the term near-bent is common).*

For $s \in \{0, 1, 2\}$, s-plateaued functions have been actively studied and have attractive much attention due to their cryptographic, algebraic, and combinatorial properties.

Two functions $f, g : \mathbb{F}_{p^n} \to \mathbb{F}_p$ are called *affinely equivalent* [18, 34] if $f(x) = ag(l(x) + b) + c$ for some linearized permutation $l(x) \in \mathbb{F}_{p^n}[x]$, $a, c \in \mathbb{F}_p$ and $b \in \mathbb{F}_{p^n}$. Note that algebraic degree, bentness of a p-ary function are affine invariants.

In this paper, we mainly focus on the secondary constructions via some known p-ary bent functions. Thus, we need to recall the following known results.

Lemma 2 (p-ary Kasami, Corollary 4 [23, 15])**.** *Let $n = 2k$ and $\lambda \in \mathbb{F}^*_{p^k}$. For any odd prime p, the p-ary monomial $f(x) = \mathrm{Tr}_1^k(\lambda x^{p^k+1})$ is a weakly regular bent function. Moreover, for $a \in \mathbb{F}_{p^n}$ the corresponding Walsh transform coefficient of $f(x)$ is equal to*

$$\widehat{\chi}_f(a) = -p^k \omega^{-\mathrm{Tr}_1^k\left(\lambda^{-1} a^{p^k+1}\right)}.$$

Lemma 3 (Sidelnikov, Corollary 3 [15])**.** *For any nonzero $\lambda \in \mathbb{F}_{p^n}$ and odd prime p, the p-ary monomial $f(x) = \mathrm{Tr}_1^n(\lambda x^2)$ is a (weakly) regular bent function. Moreover, for $a \in \mathbb{F}_{p^n}$ the corresponding Walsh transform coefficient of $f(x)$ is equal to*

$$\widehat{\chi}_f(a) = \begin{cases} \eta(\lambda)(-1)^{n-1} p^{\frac{n}{2}} \omega^{-\mathrm{Tr}_1^n(\frac{a^2}{4\lambda})}, & \text{if } p \equiv 1 \pmod{4}; \\ \eta(\lambda)(-1)^{n-1} i^n p^{\frac{n}{2}} \omega^{-\mathrm{Tr}_1^n(\frac{a^2}{4\lambda})}, & \text{if } p \equiv 3 \pmod{4}, \end{cases} \tag{2.1}$$

where $i = \sqrt{-1}$ and η is the quadratic character of $\mathbb{F}_{p^n}$.

3. QUADRATIC TERNARY BENT, NEAR-BENT AND 2-PLATEAUED FUNCTIONS FROM SOME KNOWN TERNARY BENT FUNCTIONS

In this section, we will construct several classes of quadratic ternary bent, near-bent and 2-plateaued functions. Before doing this, we need the following lemma.

Lemma 4 (Lemma 4 [33]). *Let n be a positive integer and $u, v \in \mathbb{F}_{3^n}^*$. Let $g(x)$ be a ternary function defined on $\mathbb{F}_{3^n}$. Define the ternary function $f(x)$ by*

$$f(x) = g(x) + \mathrm{Tr}_1^n(ux)\mathrm{Tr}_1^n(vx).$$

Then for any $a \in \mathbb{F}_{3^n}$, the corresponding Walsh transform coefficient of $f(x)$ is equal to

$$\begin{aligned}\widehat{\chi}_f(a) =& \frac{1}{3}[\widehat{\chi}_g(a) + \widehat{\chi}_g(a+u) + \widehat{\chi}_g(a-u)\\ &+ \widehat{\chi}_g(a-v) + \omega\widehat{\chi}_g(a-v+u) + \omega^2\widehat{\chi}_g(a-v-u)\\ &+ \widehat{\chi}_g(a+v) + \omega^2\widehat{\chi}_g(a+v+u) + \omega\widehat{\chi}_g(a+v-u)],\end{aligned} \tag{3.1}$$

where ω is a primitive 3-rd root of unity.

It was shown in [8] that quadratic monomial near-bent functions $\mathrm{Tr}_1^n(\lambda x^{p^r+1})$, $\lambda \in \mathbb{F}_{p^n}$ in odd characteristic p do not exist. Now, we can derive the following construction of ternary near-bent functions via ternary monomial bent function $\mathrm{Tr}_1^n(\lambda x^{3^r+1})$. We define a subset of $\mathbb{F}_3^3$ as

$$A = \{(0,1,1), (0,2,2), (1,1,1), (1,2,2), (2,0,1), (2,0,2), (2,1,0), (2,2,0)\}.$$

Now, using Lemmas 2 and 4, we can obtain the next theorem.

Theorem 1. *Let $k > 1$ and $n = 2k$. Let $\lambda \in \mathbb{F}_{3^k}^*$ and $u, v \in \mathbb{F}_{3^n}^*$. Let f be the ternary function defined as*

$$f(x) = \mathrm{Tr}_1^k(\lambda x^{3^k+1}) + \mathrm{Tr}_1^n(ux)\mathrm{Tr}_1^n(vx). \tag{3.2}$$

- **1):** *If $(\mathrm{Tr}_1^n(\lambda^{-1}u^{3^k}v), \mathrm{Tr}_1^k(\lambda^{-1}u^{3^k+1}), \mathrm{Tr}_1^k(\lambda^{-1}v^{3^k+1})) \in \mathbb{F}_3^3 \setminus (A \cup \{(2,0,0)\})$, then f is bent.*
- **2):** *If $(\mathrm{Tr}_1^n(\lambda^{-1}u^{3^k}v), \mathrm{Tr}_1^k(\lambda^{-1}u^{3^k+1}), \mathrm{Tr}_1^k(\lambda^{-1}v^{3^k+1})) \in A$, then f is near-bent.*
- **3):** *If $(\mathrm{Tr}_1^n(\lambda^{-1}u^{3^k}v), \mathrm{Tr}_1^k(\lambda^{-1}u^{3^k+1}), \mathrm{Tr}_1^k(\lambda^{-1}v^{3^k+1})) = (2,0,0)$, then f is a ternary 2-plateaued function. Moreover, when a runs through all elements in $\mathbb{F}_{3^n}$, the distribution of the Walsh spectrum of ternary 2-plateaued function f is given by*

$$\widehat{\chi}_f(a) = \begin{cases} 0, & \text{occurs } 3^n - 3^{n-2} \text{ times,}\\ -3^{k+1}, & \text{occurs } 3^{n-3} - 2\cdot 3^{k-2} \text{ times,}\\ -3^{k+1}\omega, & \text{occurs } 3^{n-3} + 3^{k-2} \text{ times,}\\ -3^{k+1}\omega^2, & \text{occurs } 3^{n-3} + 3^{k-2} \text{ times.}\end{cases}$$

Proof. Let $g(x) = \mathrm{Tr}_1^k(\lambda x^{3^k+1})$. For $a \in \mathbb{F}_{3^n}$, it follows from Lemma 4 that

$$\widehat{\chi}_f(a) = \triangle_1 + \triangle_3 + \triangle_3,$$

where

$$\triangle_1 = \frac{1}{3}[\widehat{\chi}_g(a) + \widehat{\chi}_g(a+u) + \widehat{\chi}_g(a-u)],$$

$$\triangle_2 = \frac{1}{3}[\widehat{\chi}_g(a-v) + \omega\widehat{\chi}_g(a-v+u) + \omega^2\widehat{\chi}_g(a-v-u)]$$

and

$$\triangle_3 = \frac{1}{3}[\widehat{\chi}_g(a+v) + \omega^2\widehat{\chi}_g(a+v+u) + \omega\widehat{\chi}_g(a+v-u)].$$

From Lemma 2, we have $\widehat{\chi}_g(a)=-3^k\omega^{-\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})}$. Then

$$\begin{aligned}\triangle_1=&\frac{1}{3}[\widehat{\chi}_g(a)+\widehat{\chi}_g(a+u)+\widehat{\chi}_g(a-u)]\\=&\frac{1}{3}[-3^k\omega^{-\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})}-3^k\omega^{-\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})-\mathrm{Tr}_1^k(\lambda^{-1}(a^{3^k}u+au^{3^k}))-\mathrm{Tr}_1^k(\lambda^{-1}u^{3^k+1})}\\&-3^k\omega^{-\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})+\mathrm{Tr}_1^k(\lambda^{-1}(a^{3^k}u+au^{3^k}))-\mathrm{Tr}_1^k(\lambda^{-1}u^{3^k+1})}]\\=&-\frac{1}{3}3^k\omega^{-\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})}[1+\omega^{-\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)-\mathrm{Tr}_1^k(\lambda^{-1}u^{3^k+1})}\\&+\omega^{\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)-\mathrm{Tr}_1^k(\lambda^{-1}u^{3^k+1})}]\end{aligned}$$

where the last identity holds due to the transitivity property of $\mathrm{Tr}_1^n(x)$, that is, $\mathrm{Tr}_1^n(x)=\mathrm{Tr}_1^k(\mathrm{Tr}_k^n(x))$.

Similarly, we have

$$\begin{aligned}\triangle_2=&\frac{1}{3}[(\widehat{\chi}_g(a-v)+\omega\widehat{\chi}_g(a-v+u)+\omega^2\widehat{\chi}_g(a-v-u)]\\=&-\frac{1}{3}3^k\omega^{-\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})+\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)-\mathrm{Tr}_1^k(\lambda^{-1}v^{3^k+1})}\\&\times[1+\omega^{1-\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)+\mathrm{Tr}_1^n(\lambda^{-1}u^{3^k}v)-\mathrm{Tr}_1^k(\lambda^{-1}u^{3^k+1})}\\&+\omega^{2+\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)-\mathrm{Tr}_1^n(\lambda^{-1}u^{3^k}v)-\mathrm{Tr}_1^k(\lambda^{-1}u^{3^k+1})}]\end{aligned}$$

and

$$\begin{aligned}\triangle_3=&\frac{1}{3}[\widehat{\chi}_g(a+v)+\omega^2\widehat{\chi}_g(a+v+u)+\omega\widehat{\chi}_g(a+v-u)]\\=&-\frac{1}{3}3^k\omega^{-\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})-\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)-\mathrm{Tr}_1^k(\lambda^{-1}v^{3^k+1})}\\&\times[1+\omega^{2-\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)-\mathrm{Tr}_1^n(\lambda^{-1}u^{3^k}v)-\mathrm{Tr}_1^k(\lambda^{-1}u^{3^k+1})}\\&+\omega^{1+\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)+\mathrm{Tr}_1^n(\lambda^{-1}u^{3^k}v)-\mathrm{Tr}_1^k(\lambda^{-1}u^{3^k+1})}].\end{aligned}$$

Let $t_0=\mathrm{Tr}_1^n(\lambda^{-1}u^{3^k}v)$, $t_1=\mathrm{Tr}_1^k(\lambda^{-1}u^{3^k+1})$ and $t_2=\mathrm{Tr}_1^k(\lambda^{-1}v^{3^k+1})$. Denote $c_1=\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)$ and $c_2=\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)$. Then the sums $\triangle_1$, $\triangle_2$ and $\triangle_3$ can be written as

$$\triangle_1=-\frac{1}{3}3^k\omega^{-\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})}(1+\omega^{-c_1-t_1}+\omega^{c_1-t_1}),\tag{3.3}$$

$$\triangle_2=-\frac{1}{3}3^k\omega^{-\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})+c_2-t_2}(1+\omega^{1-c_1+t_0-t_1}+\omega^{2+c_1-t_0-t_1})\tag{3.4}$$

and

$$\triangle_3=-\frac{1}{3}3^k\omega^{-\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})-c_2-t_2}(1+\omega^{2-c_1-t_0-t_1}+\omega^{1+c_1+t_0-t_1}).\tag{3.5}$$

Combining (3.3)-(3.5), we get

$$\begin{aligned}\widehat{\chi}_f(a)=&\triangle_1+\triangle_2+\triangle_3\\=&-\frac{1}{3}3^k\omega^{-\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})}[1+\omega^{-c_1-t_1}+\omega^{c_1-t_1}\\&+\omega^{c_2-t_2}+\omega^{-c_2-t_2}+\omega^{1-c_1+c_2+t_0-t_1-t_2}+\omega^{2+c_1+c_2-t_0-t_1-t_2}\\&+\omega^{2-c_1-c_2-t_0-t_1-t_2}+\omega^{1+c_1-c_2+t_0-t_1-t_2}].\end{aligned}\tag{3.6}$$

1) We only give the proof of the case of $(t_0, t_1, t_2) = (1, 2, 1) \in \mathbb{F}_3^3 \setminus (A \cup \{(2, 0, 0)\})$ since the others can be proven in a similar manner. Note that $(c_0, c_1) \in \mathbb{F}_3^2$ for a fixed $a \in \mathbb{F}_{3^n}$. Depending on the value of the pair (c_0, c_1) and from (3.6), we have

$$\begin{aligned}\widehat{\chi}_f(a) &= \triangle_1 + \triangle_3 + \triangle_3 \\ &= \begin{cases} 3^k \omega^{-\mathrm{Tr}_1^k(\lambda^{-1} a^{3^k+1})}, & \text{if } (c_1, c_2) = (0, 0), \\ 3^k \omega^{-\mathrm{Tr}_1^k(\lambda^{-1} a^{3^k+1})+1}, & \text{if } (c_1, c_2) \in \{(1, 0), (1, 1), (2, 0), (2, 2)\}, \\ 3^k \omega^{-\mathrm{Tr}_1^k(\lambda^{-1} a^{3^k+1})+2}, & \text{if } (c_1, c_2) \in \{(0, 1), (0, 2), (1, 2), (2, 1)\}. \end{cases}\end{aligned}$$

Hence, for all $a \in \mathbb{F}_{3^n}$, we can see that $|\widehat{\chi}_f(a)| = 3^k$ if $(t_0, t_1, t_2) = (1, 2, 1)$.

2) Let $\omega = \frac{-1+\sqrt{3}i}{2}$ where $i = \sqrt{-1}$. It is easy to verify that $\omega^2 - \omega = -\sqrt{3}i$. Similar as in 1), we only give the proof of the case of $(t_0, t_1, t_2) = (0, 1, 1) \in A$. For $a \in \mathbb{F}_{3^n}$, from (3.6), we have

$$\begin{aligned}\widehat{\chi}_f(a) &= \triangle_1 + \triangle_3 + \triangle_3 \\ &= \begin{cases} 3^{k+1/2} i \omega^{-\mathrm{Tr}_1^k(\lambda^{-1} a^{3^k+1})}, & \text{if } (c_1, c_2) = (0, 0), \\ 3^{k+1/2} i \omega^{-\mathrm{Tr}_1^k(\lambda^{-1} a^{3^k+1})+1}, & \text{if } (c_1, c_2) \in \{(1, 2), (2, 1)\}, \\ 0, & \text{otherwise}. \end{cases}\end{aligned}$$

Consequently, for $a \in \mathbb{F}_{3^n}$, $|\widehat{\chi}_f(a)| \in \{0, 3^{k+1/2}\}$ if $(t_0, t_1, t_2) = (0, 1, 1)$.

3) If $t_0 = 2$, $t_1 = 0$ and $t_2 = 0$, by (3.3)-(3.5), we have

$$\begin{aligned}\widehat{\chi}_f(a) &= \triangle_1 + \triangle_3 + \triangle_3 \\ &= -\frac{1}{3} 3^k \omega^{-\mathrm{Tr}_1^k(\lambda^{-1} a^{3^k+1})} (1 + \omega^{-c_1} + \omega^{c_1})(1 + \omega^{c_2} + \omega^{-c_2}) \\ &= \begin{cases} -3^{k+1} \omega^{-\mathrm{Tr}_1^k(\lambda^{-1} a^{3^k+1})}, & \text{if } (c_1, c_2) = (0, 0), \\ 0, & \text{otherwise}. \end{cases} \qquad (3.7)\end{aligned}$$

Therefore, $|\widehat{\chi}_f(a)| \in \{0, 3^{k+1}\}$ for all $a \in \mathbb{F}_{3^n}$. Then f is 2-plateaued.

By a complex and lengthy computation, we can obtain the value distribution of the Walsh transform of 2-plateaued function f, and the details are presented in Appendix 6. □

Example 1. *Let $k = 3$, $n = 6$. Let α be the generator of $\mathbb{F}_{3^6}^*$ with $\alpha^6 - \alpha^4 + \alpha^2 - \alpha - 2 = 0$.*

1): *Take $\lambda = \alpha^{84}$, $u = \alpha^4$ and $v = \alpha^6$. Then f defined by (3.2) is*

$$f(x) = \mathrm{Tr}_1^3(\alpha^{84} x^{28}) + \mathrm{Tr}_1^6(\alpha^4 x)\mathrm{Tr}_1^6(\alpha^6 x).$$

By a Magma program, we can see that $\mathrm{Tr}_1^6(\lambda^{-1} u^{27} v) = 1$, $\mathrm{Tr}_1^3(\lambda^{-1} u^{28}) = 0$, $\mathrm{Tr}_1^3(\lambda^{-1} v^{28}) = 0$ and f is bent.

2): *Take $\lambda = \alpha^{84}$, $u = \alpha^7$ and $v = \alpha^{25}$. Then f defined by (3.2) is*

$$f(x) = \mathrm{Tr}_1^3(\alpha^{84} x^{28}) + \mathrm{Tr}_1^6(\alpha^7 x)\mathrm{Tr}_1^6(\alpha^{25} x).$$

By a Magma program, we can see that $\mathrm{Tr}_1^6(\lambda^{-1} u^{27} v) = 2$, $\mathrm{Tr}_1^3(\lambda^{-1} u^{28}) = 2$, $\mathrm{Tr}_1^3(\lambda^{-1} v^{28}) = 0$ and f is near-bent.

3): *Take $\lambda = \alpha^{84}$, $u = \alpha^4$ and $v = \alpha^{25}$. Then f defined by (3.2) is*

$$f(x) = \mathrm{Tr}_1^3(\alpha^{84} x^{28}) + \mathrm{Tr}_1^6(\alpha^4 x)\mathrm{Tr}_1^6(\alpha^{25} x).$$

By a Magma program, we can see that $\mathrm{Tr}_1^6(\lambda^{-1} u^{27} v) = 2$, $\mathrm{Tr}_1^3(\lambda^{-1} u^{28}) = 0$, $\mathrm{Tr}_1^3(\lambda^{-1} v^{28}) = 0$ and f is 2-plateaued. Moreover, the value distribution of the

Walsh transform of f is

$$\widehat{\chi}_f(a)=\begin{cases}0, & \textit{occurs } 648 \textit{ times,}\\ -81, & \textit{occurs } 21 \textit{ times,}\\ -81\omega, & \textit{occurs } 30 \textit{ times,}\\ -81\omega^2, & \textit{occurs } 30 \textit{ times.}\end{cases}$$

Thus, our computer experiments are consistent with the results given in Theorem 1.

We define a subset of $\mathbb{F}_3^3$ as

$$B=\{(0,1,1),(0,2,2),(1,0,1),(1,0,2),(1,1,0),(1,2,0),(2,1,1),(2,2,2)\}.$$

Similarly, by Lemmas 3 and 4, we can prove the following theorem.

Theorem 2. *Let n be a positive integer with $n>3$ and $\lambda,u,v\in\mathbb{F}_{3^n}^*$. The ternary function f defined as*

$$f(x)=\mathrm{Tr}_1^n(\lambda x^2)+\mathrm{Tr}_1^n(ux)\mathrm{Tr}_1^n(vx). \tag{3.8}$$

1): *If $(\mathrm{Tr}_1^n(\lambda^{-1}uv),\mathrm{Tr}_1^n(\lambda^{-1}u^2),\mathrm{Tr}_1^n(\lambda^{-1}v^2))\in\mathbb{F}_3^3\setminus(B\cup\{(1,0,0)\})$, then f is bent.*

2): *If $(\mathrm{Tr}_1^n(\lambda^{-1}uv),\mathrm{Tr}_1^n(\lambda^{-1}u^2),\mathrm{Tr}_1^n(\lambda^{-1}v^2))\in B$, then f is near-bent.*

3): *If $(\mathrm{Tr}_1^n(\lambda^{-1}uv),\mathrm{Tr}_1^n(\lambda^{-1}u^2),\mathrm{Tr}_1^n(\lambda^{-1}v^2))=(1,0,0)$, then f is a ternary 2-plateaued function. Moreover, when a runs through all elements in $\mathbb{F}_{3^n}$, the distribution of the Walsh spectrum of ternary 2-plateaued function f is shown as follows.*

i) For the case n is odd,

$$\widehat{\chi}_f(a)=\begin{cases}0, & \textit{occurs } 3^n-3^{n-2} \textit{ times,}\\ \eta(\lambda)i^n3^{\frac{n}{2}+1}, & \textit{occurs } 3^{n-3} \textit{ times,}\\ \eta(\lambda)i^n3^{\frac{n}{2}+1}\omega, & \textit{occurs } 3^{n-3}+\eta(\lambda)i^{n+1}3^{\frac{n-3}{2}} \textit{ times,}\\ \eta(\lambda)i^n3^{\frac{n}{2}+1}\omega^2, & \textit{occurs } 3^{n-3}-\eta(\lambda)i^{n+1}3^{\frac{n-3}{2}} \textit{ times.}\end{cases}$$

i) For the case n is even,

$$\widehat{\chi}_f(a)=\begin{cases}0, & \textit{occurs } 3^n-3^{n-2} \textit{ times,}\\ -\eta(\lambda)i^n3^{\frac{n}{2}+1}, & \textit{occurs } 3^{n-3}-2\eta(\lambda)i^n3^{\frac{n}{2}-2} \textit{ times,}\\ -\eta(\lambda)i^n3^{\frac{n}{2}+1}\omega, & \textit{occurs } 3^{n-3}+\eta(\lambda)i^n3^{\frac{n}{2}-2} \textit{ times,}\\ -\eta(\lambda)i^n3^{\frac{n}{2}+1}\omega^2, & \textit{occurs } 3^{n-3}+\eta(\lambda)i^n3^{\frac{n}{2}-2} \textit{ times.}\end{cases}$$

Proof. The proof is similar to that of Theorem 1 and is omitted. □

Remark 1. *By Theorems 1 and 2, one can conclude that a ternary function given by adding the product of the product of two linear functions to known ternary bent function $\mathrm{Tr}_1^k(\lambda x^{3^k+1})$ or $\mathrm{Tr}_1^n(\lambda x^2)$ must be among ternary bent, near-bent and 2-plateaued function. It is interesting to find other ternary bent functions $f(x)$ such that $f(x)+\mathrm{Tr}_1^n(ux)\mathrm{Tr}_1^n(vx)$ are also bent.*

Example 2. *Let $n=4$ and let α be the generator of $\mathbb{F}_{3^4}^*$ with $\alpha^4-\alpha^3-1=0$.*

1): *Take $\lambda=\alpha$, $u=\alpha^4$ and $v=\alpha^7$. Then f defined by (3.8) is*

$$f(x)=\mathrm{Tr}_1^4(\alpha x^2)+\mathrm{Tr}_1^4(\alpha^4x)\mathrm{Tr}_1^4(\alpha^7x).$$

By a Magma program, we can see that $\mathrm{Tr}_1^4(\lambda^{-1}uv)=2,\mathrm{Tr}_1^4(\lambda^{-1}u^2)=2,\mathrm{Tr}_1^4(\lambda^{-1}v^2)=0$ and f is bent. This is consistent with the results given in Theorem 2.

2): *Take $\lambda=\alpha$, $u=\alpha^4$ and $v=\alpha^8$. Then f defined by (3.8) is*

$$f(x)=\mathrm{Tr}_1^4(\alpha x^2)+\mathrm{Tr}_1^4(\alpha^4x)\mathrm{Tr}_1^4(\alpha^8x).$$

By a Magma program, we can see that $\mathrm{Tr}_1^4(\lambda^{-1}uv)=1,\mathrm{Tr}_1^4(\lambda^{-1}u^2)=2,\mathrm{Tr}_1^4(\lambda^{-1}v^2)=0$ and f is near-bent.

3): *Take $\lambda=\alpha$, $u=\alpha^{16}$ and $v=\alpha^{8}$. Then f defined by (3.8) is*

$$f(x)=\mathrm{Tr}_1^4(\alpha x^2)+\mathrm{Tr}_1^4(\alpha^{16}x)\mathrm{Tr}_1^4(\alpha^8 x).$$

By a Magma program, we can see that $\mathrm{Tr}_1^4(\lambda^{-1}uv)=1$, $\mathrm{Tr}_1^4(\lambda^{-1}u^2)=0$, $\mathrm{Tr}_1^4(\lambda^{-1}v^2)=0$ and f is 2-plateaued. The the value distribution of the Walsh transform of f is

$$\widehat{\chi}_f(a)=\begin{cases}0, & \text{occurs } 72 \text{ times,}\\ 27, & \text{occurs } 5 \text{ times,}\\ 27\omega, & \text{occurs } 2 \text{ times,}\\ 27\omega^2, & \text{occurs } 2 \text{ times.}\end{cases}$$

Thus, our computer experiments are consistent with the results given in Theorem 2.

Below, we identify $\mathbb{F}_{3^n}$ (where $n=2k$) with $\mathbb{F}_{3^k}\times\mathbb{F}_{3^k}$. For $a=(a_1,a_2),b=(b_1,b_2)\in\mathbb{F}_{3^k}\times\mathbb{F}_{3^k}$, the scalar product in $\mathbb{F}_{3^n}$ can be defined as $\langle(a_1,a_2),(b_1,b_2)\rangle=\mathrm{Tr}_1^k(a_1b_1+a_2b_2)$. The well-known Maiorana-McFarland class of ternary bent functions can be defined as

$$g(x,y)=\mathrm{Tr}_1^k(x\pi(y))+h(y),(x,y)\in\mathbb{F}_{3^k}\times\mathbb{F}_{3^k}$$

where $\pi:\mathbb{F}_{3^k}\to\mathbb{F}_{3^k}$ is a permutation and h is a ternary function over $\mathbb{F}_{3^k}$, and its dual is given by

$$g^*(x,y)=-\mathrm{Tr}_1^k(y\pi^{-1}(x))+h(\pi^{-1}(x))$$

where π^{-1} denotes the inverse mapping of the permutation π [20]. This together with the definition of the dual function implies that for each $a=(a_1,a_2)\in\mathbb{F}_{3^n}$

$$\widehat{\chi}_g(a_1,a_2)=3^k\omega^{-\mathrm{Tr}_1^k(a_2\pi^{-1}(a_1))+h(\pi^{-1}(a_1))}. \tag{3.9}$$

Similar to Lemma 4, we have the following result.

Lemma 5. *Let $n=2k$ be a even positive integer and $(u_1,u_2),(v_1,v_2)\in\mathbb{F}_{3^k}\times\mathbb{F}_{3^k}$. Let $g(x,y)$ be a ternary function defined on $\mathbb{F}_{3^k}\times\mathbb{F}_{3^k}$. Let $f(x,y)$ be the ternary function defined as*

$$f(x,y)=g(x,y)+\mathrm{Tr}_1^k(u_1x+u_2y)\mathrm{Tr}_1^k(v_1x+v_2y).$$

Then for any $(a_1,a_2)\in\mathbb{F}_{3^k}\times\mathbb{F}_{3^k}$, the corresponding Walsh transform coefficient of $f(x,y)$ is equal to

$$\begin{aligned}\widehat{\chi}_f(a_1,a_2)=&\frac{1}{3}[\widehat{\chi}_g(a_1,a_2)+\widehat{\chi}_g(a_1+u_1,a_2+u_2)+\widehat{\chi}_g(a_1-u_1,a_2-u_2)\\&+\widehat{\chi}_g(a_1-v_1,a_2-v_2)+\omega\widehat{\chi}_g(a_1-v_1+u_1,a_2-v_2+u_2)\\&+\omega^2\widehat{\chi}_g(a_1-v_1-u_1,a_2-v_2-u_2)+\widehat{\chi}_g(a_1+v_1,a_2+v_2)\\&+\omega^2\widehat{\chi}_g(a_1+v_1+u_1,a_2+v_2+u_2)+\omega\widehat{\chi}_g(a_1+v_1-u_1,a_2+v_2-u_2)]\end{aligned}$$

where ω is a primitive 3-rd root of unity.

The following theorem will employ the linearized permutation polynomial over $\mathbb{F}_{3^k}$ to give new ternary bent functions from the class of Maiorana-McFarland.

Theorem 3. *Let A be the subset of $\mathbb{F}_3^3$ defined as above. Let $n=2k$ and $u=(u_1,u_2),v=(v_1,v_2)$ be two nonzero elements in $\mathbb{F}_{3^k}\times\mathbb{F}_{3^k}$. Assume that π is a linearized permutation polynomial over $\mathbb{F}_{3^k}$. Let $f(x,y)$ be the ternary function given by*

$$f(x,y)=\mathrm{Tr}_1^k(x\pi(y))+\mathrm{Tr}_1^k(y)+\mathrm{Tr}_1^k(u_1x+u_2y)\mathrm{Tr}_1^k(v_1x+v_2y). \tag{3.10}$$

- **1):** *If $\left(\mathrm{Tr}_1^k(u_2\pi^{-1}(v_1)+v_2\pi^{-1}(u_1)),\mathrm{Tr}_1^k(u_2\pi^{-1}(u_1)),\mathrm{Tr}_1^k(v_2\pi^{-1}(v_1))\right)\in\mathbb{F}_3^3\setminus(A\cup\{(2,0,0)\})$, then $f(x,y)$ is bent.*
- **2):** *If $\left(\mathrm{Tr}_1^k(u_2\pi^{-1}(v_1)+v_2\pi^{-1}(u_1)),\mathrm{Tr}_1^k(u_2\pi^{-1}(u_1)),\mathrm{Tr}_1^k(v_2\pi^{-1}(v_1))\right)\in A$, then $f(x,y)$ is near-bent.*

3): *If* $\big(\mathrm{Tr}_1^k(u_2\pi^{-1}(v_1)+v_2\pi^{-1}(u_1)), \mathrm{Tr}_1^k(u_2\pi^{-1}(u_1)), \mathrm{Tr}_1^k(v_2\pi^{-1}(v_1))\big) = (2,0,0)$, *then* $f(x,y)$ *is a ternary 2-plateaued function.*

Proof. Let $g(x,y) = \mathrm{Tr}_1^k(x\pi(y)) + \mathrm{Tr}_1^k(y)$. Applying Lemma 5, for each $(a_1,a_2) \in \mathbb{F}_{3^k} \times \mathbb{F}_{3^k}$, we have

$$\widehat{\chi}_f(a_1,a_2) = \triangle_1 + \triangle_2 + \triangle_3,$$

where

$$\triangle_1 = \frac{1}{3}[\widehat{\chi}_g(a_1,a_2) + \widehat{\chi}_g(a_1+u_1,a_2+u_2) + \widehat{\chi}_g(a_1-u_1,a_2-u_2)],$$

$$\begin{aligned}\triangle_2 =& \frac{1}{3}[\widehat{\chi}_g(a_1-v_1,a_2-v_2) + \omega\widehat{\chi}_g(a_1-v_1+u_1,a_2-v_2+u_2)\\ &+ \omega^2\widehat{\chi}_g(a_1-v_1-u_1,a_2-v_2-u_2)].\end{aligned}$$

and

$$\begin{aligned}\triangle_3 =& \frac{1}{3}[\widehat{\chi}_g(a_1+v_1,a_2+v_2) + \omega^2\widehat{\chi}_g(a_1+v_1+u_1,a_2+v_2+u_2)\\ &+ \omega\widehat{\chi}_g(a_1+v_1-u_1,a_2+v_2-u_2)].\end{aligned}$$

Note that π is a linearized permutation polynomial, and thus, π^{-1} is also a linearized permutation polynomial. From (3.9), we can compute the sums $\triangle_1$, $\triangle_2$ and $\triangle_3$ respectively.

$$\begin{aligned}\triangle_1 =& \frac{1}{3}\widehat{\chi}_g(a_1,a_2)[1 + \omega^{-\mathrm{Tr}_1^k(a_2\pi^{-1}(u_1))-\mathrm{Tr}_1^k(u_2\pi^{-1}(a_1))+\mathrm{Tr}_1^k(\pi^{-1}(u_1))-\mathrm{Tr}_1^k(u_2\pi^{-1}(u_1))}\\ &+ \omega^{\mathrm{Tr}_1^k(a_2\pi^{-1}(u_1))+\mathrm{Tr}_1^k(u_2\pi^{-1}(a_1))-\mathrm{Tr}_1^k(\pi^{-1}(u_1))-\mathrm{Tr}_1^k(u_2\pi^{-1}(u_1))}]\end{aligned}\tag{3.11}$$

Similarly, we have

$$\begin{aligned}\triangle_2 =& \frac{1}{3}\widehat{\chi}_g(a_1,a_2)\omega^{\mathrm{Tr}_1^k(a_2\pi^{-1}(v_1))+\mathrm{Tr}_1^k(v_2\pi^{-1}(a_1))-\mathrm{Tr}_1^k(\pi^{-1}(v_1))-\mathrm{Tr}_1^k(v_2\pi^{-1}(v_1))}\\ &\times [1 + \omega^{1-\mathrm{Tr}_1^k(a_2\pi^{-1}(u_1))-\mathrm{Tr}_1^k(u_2\pi^{-1}(a_1))+\mathrm{Tr}_1^k(\pi^{-1}(u_1))-\mathrm{Tr}_1^k(u_2\pi^{-1}(u_1))}\\ &\times \omega^{\mathrm{Tr}_1^k(u_2\pi^{-1}(v_1))+\mathrm{Tr}_1^k(v_2\pi^{-1}(u_1))}\\ &+ \omega^{2+\mathrm{Tr}_1^k(a_2\pi^{-1}(u_1))+\mathrm{Tr}_1^k(u_2\pi^{-1}(a_1))-\mathrm{Tr}_1^k(\pi^{-1}(u_1))-\mathrm{Tr}_1^k(u_2\pi^{-1}(u_1))}\\ &\times \omega^{-\mathrm{Tr}_1^k(u_2\pi^{-1}(v_1))-\mathrm{Tr}_1^k(v_2\pi^{-1}(u_1))}]\end{aligned}\tag{3.12}$$

and

$$\begin{aligned}\triangle_3 =& \frac{1}{3}\widehat{\chi}_g(a_1,a_2)\omega^{-\mathrm{Tr}_1^k(a_2\pi^{-1}(v_1))-\mathrm{Tr}_1^k(v_2\pi^{-1}(a_1))+\mathrm{Tr}_1^k(\pi^{-1}(v_1))-\mathrm{Tr}_1^k(v_2\pi^{-1}(v_1))}\\ &\times [1 + \omega^{2-\mathrm{Tr}_1^k(a_2\pi^{-1}(u_1))-\mathrm{Tr}_1^k(u_2\pi^{-1}(a_1))+\mathrm{Tr}_1^k(\pi^{-1}(u_1))-\mathrm{Tr}_1^k(u_2\pi^{-1}(u_1))}\\ &\times \omega^{-\mathrm{Tr}_1^k(u_2\pi^{-1}(v_1))-\mathrm{Tr}_1^k(v_2\pi^{-1}(u_1))}\\ &+ \omega^{1+\mathrm{Tr}_1^k(a_2\pi^{-1}(u_1))+\mathrm{Tr}_1^k(u_2\pi^{-1}(a_1))-\mathrm{Tr}_1^k(\pi^{-1}(u_1))-\mathrm{Tr}_1^k(u_2\pi^{-1}(u_1))}\\ &\times \omega^{\mathrm{Tr}_1^k(u_2\pi^{-1}(v_1))+\mathrm{Tr}_1^k(v_2\pi^{-1}(u_1))}].\end{aligned}\tag{3.13}$$

For convenience of presentation, we let $t_0 = \mathrm{Tr}_1^k(u_2\pi^{-1}(v_1)) + \mathrm{Tr}_1^k(v_2\pi^{-1}(u_1))$, $t_1 = \mathrm{Tr}_1^k(u_2\pi^{-1}(u_1))$ and $t_2 = \mathrm{Tr}_1^k(v_2\pi^{-1}(v_1))$ and denote $c_1 = \mathrm{Tr}_1^k(a_2\pi^{-1}(u_1)) + \mathrm{Tr}_1^k(u_2\pi^{-1}(a_1)) - \mathrm{Tr}_1^k(\pi^{-1}(u_1))$ and $c_2 = \mathrm{Tr}_1^k(a_2\pi^{-1}(v_1)) + \mathrm{Tr}_1^k(v_2\pi^{-1}(a_1)) - \mathrm{Tr}_1^k(\pi^{-1}(v_1))$. Thus, (3.11), (3.12) and (3.13) can be written as

$$\triangle_1 = \tfrac{1}{3}\widehat{\chi}_g(a_1,a_2)\left(1 + \omega^{-c_1-t_1} + \omega^{c_1-t_1}\right),$$

$$\triangle_2 =\frac{1}{3}\widehat{\chi}_g(a_1,a_2))\omega^{c_2-t_2}(1+\omega^{1-c_1+t_0-t_1}+\omega^{2+c_1-t_0-t_1})$$

and

$$\triangle_3 =\frac{1}{3}\widehat{\chi}_g(a_1,a_2))\omega^{-c_2-t_2}(1+\omega^{2-c_1-t_0-t_1}+\omega^{1+c_1+t_0-t_1}),$$

Similar as the proof of Theorem 1, detailed discussing of the possible values of t_i and c_i leads to the desired conclusion. □

Remark 2. *To obtain our constructions in Theorem 3, we need to know the compositional inverse of a given linearized permutation polynomial over* $\mathbb{F}_{3^k}$*. It is clear that the simplest suitable linearized permutation polynomial* π *over* $\mathbb{F}_{3^k}$ *in Theorem 3 is* y^{3^i} *where* $0\le i\le n-1$*. In addition, the linearized permutation binomials in Lemma 1 with* $p=3$ *can also be employed to construct ternary functions in Theorem 3.*

Example 3. *Let* $k=4$ *and* $n=8$*. Let* α *be the generator of* $\mathbb{F}_{3^4}^*$ *with* $\alpha^4-\alpha^3-1=0$*. Take* $\pi(y)=y^9+\alpha y$*, by Lemma 1, we can get* $\pi^{-1}(y)=(\alpha^{10}-1)^{-1}(\alpha^9 y-y^9)$*.*

1): *Let* $u=(u_1,u_2)=(\alpha^4,\alpha^5)$ *and* $v=(v_1,v_2)=(\alpha^{10},\alpha^2)$*. Then* $f(x,y)$ *defined by (3.10) is*

$$f(x,y)=\mathrm{Tr}_1^4(x(y^9+\alpha y))+\mathrm{Tr}_1^4(y)+\mathrm{Tr}_1^4(\alpha^4x+\alpha^5y)\mathrm{Tr}_1^4(\alpha^{10}x+\alpha^2y).$$

Using Magma, we can verify that $\mathrm{Tr}_1^4(u_2\pi^{-1}(v_1))+\mathrm{Tr}_1^k(v_2\pi^{-1}(u_1))=1$, $\mathrm{Tr}_1^4(u_2\pi^{-1}(u_1)))=0$, $\mathrm{Tr}_1^4(v_2\pi^{-1}(v_1)))=0$ *and* $f(x,y)$ *is bent.*

2): *Let* $u=(u_1,u_2)=(\alpha^{10},\alpha^{11})$ *and* $v=(v_1,v_2)=(\alpha^{10},\alpha^{73})$*. Then* $f(x,y)$ *defined by (3.10) is*

$$f(x,y)=\mathrm{Tr}_1^4(x(y^9+\alpha y))+\mathrm{Tr}_1^4(y)+\mathrm{Tr}_1^4(\alpha^{10}x+\alpha^{11}y)\mathrm{Tr}_1^4(\alpha^{10}x+\alpha^{73}y).$$

Using Magma, we can verify that $\mathrm{Tr}_1^4(u_2\pi^{-1}(v_1))+\mathrm{Tr}_1^4(v_2\pi^{-1}(u_1))=2$, $\mathrm{Tr}_1^4(u_2\pi^{-1}(u_1)))=2$, $\mathrm{Tr}_1^4(v_2\pi^{-1}(v_1)))=0$ *and* $f(x,y)$ *is near-bent.*

3): *Let* $u=(u_1,u_2)=(\alpha^4,\alpha^5)$ *and* $v=(v_1,v_2)=(\alpha^{10},\alpha^{46})$*. Then* $f(x,y)$ *defined by (3.10) is*

$$f(x,y)=\mathrm{Tr}_1^4(x(y^9+\alpha y))+\mathrm{Tr}_1^4(y)+\mathrm{Tr}_1^4(\alpha^4x+\alpha^5y)\mathrm{Tr}_1^4(\alpha^{10}x+\alpha^{46}y).$$

Using Magma, we can verify that $\mathrm{Tr}_1^4(u_2\pi^{-1}(v_1))+\mathrm{Tr}_1^4(v_2\pi^{-1}(u_1))=2$, $\mathrm{Tr}_1^4(u_2\pi^{-1}(u_1)))=0$, $\mathrm{Tr}_1^4(v_2\pi^{-1}(v_1)))=0$ *and* $f(x,y)$ *is 2-plateaued.*

Our computer experiments are consistent with the results given in Theorem 3.

4. Two new class of non-quadratic p-ary bent functions of algebraic degree p

In [33], the authors provided a class of non-quadratic p-ary bent functions $f(x)=\mathrm{Tr}_1^k(\lambda x^{p^i+1})+c\mathrm{Tr}_1^n(x)^{p-1}$ of the algebraic degree $p-1$, where $\frac{n}{\gcd(i,n)}$ is odd. In this section, two new class of non-quadratic p-ary bent functions of algebraic degree p will be obtained. Before doing this, we need the following lemma.

Lemma 6. *Let* p *be an odd prime and* n *be a positive integer. Let* $g(x)$ *be a* p*-ary function defined on* $\mathbb{F}_{p^n}$*. Let* $f(x)$ *be the* p*-ary function defined by*

$$f(x)=g(x)-\mathrm{Tr}_1^n(ux)\mathrm{Tr}_1^n(x)^{p-1},$$

where $c \in \mathbb{F}_p^*$ *and* $u \in \mathbb{F}_{p^n}^*$ *. Then, for every* $a \in \mathbb{F}_{p^n}$ *the corresponding Walsh transform coefficient of* $f(x)$ *is equal to*

$$\widehat{\chi}_f(a) \;=\; \frac{1}{p}\sum_{j=0}^{p-1}\widehat{\chi}_g(a+j) - \frac{1}{p}\sum_{j=0}^{p-1}\widehat{\chi}_g(a+u+j) + \widehat{\chi}_g(a+u),$$

where ω is a primitive p-th root of unity.

Proof. For $i = 0, 1, \cdots, p-1$, we define $T_i = \{x \in \mathbb{F}_{p^n} | \mathrm{Tr}_1^n(x) = i\}$. To calculate the Walsh transform coefficient of $f(x)$ evaluated at $a \in \mathbb{F}_{p^n}$, we define the following exponential sums

$$S_i(a) = \sum_{x\in T_i}\omega^{g(x)-\mathrm{Tr}_1^n(ax)}, Q_i(a+u) = \sum_{x\in T_i}\omega^{g(x)-\mathrm{Tr}_1^n((a+u)x)},$$

where $i = 0, 1, \cdots, p-1$. Then we have

$$\begin{aligned}
&\widehat{\chi}_f(a)\\
=&\sum_{x\in\mathbb{F}_{p^n}}\omega^{f(x)-\mathrm{Tr}_1^n(ax)}\\
=&\sum_{x\in\mathbb{F}_{p^n}}\omega^{g(x)-\mathrm{Tr}_1^n(ux)\mathrm{Tr}_1^n(x)^{p-1}-\mathrm{Tr}_1^n(ax)}\\
=&\sum_{x\in T_0}\omega^{g(x)-\mathrm{Tr}_1^n(ax)} + \sum_{x\in T_1}\omega^{g(x)-\mathrm{Tr}_1^n((a+u)x)} + \cdots + \sum_{x\in T_{p-1}}\omega^{g(x)-\mathrm{Tr}_1^n((a+u)x)}\\
=&S_0(a) - Q_0(a+u) + \widehat{\chi}_g(a+u).
\end{aligned}$$

For a $j \in \mathbb{F}_p$, we can derive

$$\sum_{x\in T_i}\omega^{g(x)-\mathrm{Tr}_1^n((a+j)x)} = \sum_{x\in T_i}\omega^{g(x)-\mathrm{Tr}_1^n(ax)-j\mathrm{Tr}_1^n(x)} = \omega^{-ji}S_i(a). \tag{4.1}$$

It then follows that $\widehat{\chi}_g(a+j) = \sum_{i=0}^{p-1}\omega^{-ji}S_i(a)$. For $j \in \{0, 1, \cdots, p-1\}$, we can get

$$\begin{cases}
\widehat{\chi}_g(a) = \sum_{i=0}^{p-1}S_i(a)\\
\widehat{\chi}_g(a+1) = \sum_{i=0}^{p-1}\omega^{-i}S_i(a)\\
\cdots\\
\widehat{\chi}_g(a+p-1) = \sum_{i=0}^{p-1}\omega^{-(p-1)i}S_i(a)
\end{cases}. \tag{4.2}$$

Note that $1 + \omega + \cdots + \omega^{p-1} = 0$. Adding p equations of (4.2) gives

$$S_0(a) = \frac{1}{p}\sum_{j=0}^{p-1}\widehat{\chi}_g(a+j).$$

Substituting a by $a+u$ in (4.1) and repeating above process, we have

$$Q_0(a+u) = \frac{1}{p}\sum_{j=0}^{p-1}\widehat{\chi}_g(a+u+j).$$

Finally, we have

$$\widehat{\chi}_f(a) \;=\; \frac{1}{p}\sum_{j=0}^{p-1}\widehat{\chi}_g(a+j) - \frac{1}{p}\sum_{j=0}^{p-1}\widehat{\chi}_g(a+u+j) + \widehat{\chi}_g(a+u).$$

□

Theorem 4. *Let $n = 2k$ with $k > 1$ and $\lambda \in \mathbb{F}_{p^k}^*$, $u \in \mathbb{F}_{p^n} \setminus \mathbb{F}_p$ such that $\mathrm{Tr}_1^k(\lambda^{-1}) = 0$ and $\mathrm{Tr}_1^n(\lambda^{-1}u) = 0$. Then the p-ary function*

$$f(x) = \mathrm{Tr}_1^k(\lambda x^{p^k+1}) - \mathrm{Tr}_1^n(ux)\mathrm{Tr}_1^n(x)^{p-1} \tag{4.3}$$

is a weakly regular bent function. Moreover, for $a \in \mathbb{F}_{p^n}$ the corresponding Walsh transform coefficient of $f(x)$ is equal to

$$\widehat{\chi}_f(a) = \begin{cases} -p^k\omega^{-\mathrm{Tr}_1^k(\lambda^{-1}a^{p^k+1})}, & \text{if } \mathrm{Tr}_1^n(\lambda^{-1}a) = 0\ ; \\ -p^k\omega^{-\mathrm{Tr}_1^k(\lambda^{-1}(a+u)^{p^k+1})}, & \text{if } \mathrm{Tr}_1^n(\lambda^{-1}a) \neq 0. \end{cases}$$

Proof. Let $g(x) = \mathrm{Tr}_1^k(\lambda x^{p^k+1})$. By lemma 2, $g(x)$ is a p-ary bent function and

$$\widehat{\chi}_g(a) = -p^k\omega^{-\mathrm{Tr}_1^k(\lambda^{-1}a^{p^k+1})}$$

for $a \in \mathbb{F}_{p^n}$. It then follows from Lemma 6 that

$$\begin{aligned}
\widehat{\chi}_f(a) =& \frac{1}{p}\sum_{j=0}^{p-1}\widehat{\chi}_g(a+j) - \frac{1}{p}\sum_{j=0}^{p-1}\widehat{\chi}_g(a+u+j) + \widehat{\chi}_g(a+u) \\
=& -p^k\Bigg(\frac{1}{p}\sum_{j=0}^{p-1}\omega^{-\mathrm{Tr}_1^k(\lambda^{-1}a^{p^k+1})-j\mathrm{Tr}_1^k(\lambda^{-1}(a^{p^k}+a))-j^2\mathrm{Tr}_1^k(\lambda^{-1})} \\
&- \frac{1}{p}\sum_{j=0}^{p-1}\omega^{-\mathrm{Tr}_1^k(\lambda^{-1}(a+u)^{p^k+1})-j\mathrm{Tr}_1^k(\lambda^{-1}((a+u)^{p^k}+a+u))-j^2\mathrm{Tr}_1^k(\lambda^{-1})} \\
&+ \omega^{-\mathrm{Tr}_1^k(\lambda^{-1}(a+u)^{p^k+1})}\Bigg) \\
=& -p^k\Bigg(\frac{1}{p}\sum_{j=0}^{p-1}\omega^{-\mathrm{Tr}_1^k(\lambda^{-1}a^{p^k+1})-j\mathrm{Tr}_1^n(\lambda^{-1}a)-j^2\mathrm{Tr}_1^k(\lambda^{-1})} \\
&- \frac{1}{p}\sum_{j=0}^{p-1}\omega^{-\mathrm{Tr}_1^k(\lambda^{-1}(a+u)^{p^k+1})-j\mathrm{Tr}_1^n(\lambda^{-1}a)-j\mathrm{Tr}_1^n(\lambda^{-1}u)-j^2\mathrm{Tr}_1^k(\lambda^{-1})} \\
&+ \omega^{-\mathrm{Tr}_1^k(\lambda^{-1}(a+u)^{p^k+1})}\Bigg)
\end{aligned} \tag{4.4}$$

Since $\mathrm{Tr}_1^k(\lambda^{-1}) = 0$ and $\mathrm{Tr}_1^n(\lambda^{-1}u) = 0$, (4.4) can be simplified to

$$\begin{aligned}
\widehat{\chi}_f(a) =& -p^k\Bigg(\omega^{-\mathrm{Tr}_1^k(\lambda^{-1}a^{p^k+1})}\frac{1}{p}\sum_{j=0}^{p-1}\omega^{-j\mathrm{Tr}_1^n(\lambda^{-1}a)} \\
&- \omega^{-\mathrm{Tr}_1^k(\lambda^{-1}(a+u)^{p^k+1})}\big(\frac{1}{p}\sum_{j=0}^{p-1}\omega^{-j\mathrm{Tr}_1^n(\lambda^{-1}a)} - 1\big)\Bigg)
\end{aligned}$$

Clearly, if $\mathrm{Tr}_1^n(\lambda^{-1}a) = 0$, then

$$\widehat{\chi}_f(a) = -p^k\omega^{-\mathrm{Tr}_1^k(\lambda^{-1}a^{p^k+1})}.$$

If $\mathrm{Tr}_1^n(\lambda^{-1}a) \neq 0$, then

$$\widehat{\chi}_f(a) = -p^k\omega^{-\mathrm{Tr}_1^k(\lambda^{-1}(a+u)^{p^k+1})}.$$

Therefore $|\widehat{\chi}_f(a)| = p^k$ for all $a \in \mathbb{F}_{p^n}$. $\square$

Remark 3. *1) If $k = 1$, i.e., $n = 2$, then $\mathrm{Tr}_1^k(\lambda^{-1}) = \lambda^{-1} \neq 0$ for $\lambda \in \mathbb{F}_p^*$. It is clear that $\mathrm{Tr}_1^n(ux)\mathrm{Tr}_1^n(x)^{p-1} = u\mathrm{Tr}_1^n(x)$ if $u \in \mathbb{F}_p^*$.*

2) It was shown in [17] that the algebraic degree of p-ary weakly bent functions with $(p-1)n \geq 4$ over $\mathbb{F}_{p^n}$ is at most $\frac{(p-1)n}{2}$. Clearly, the algebraic degree of $\mathrm{Tr}_1^n(ux)\mathrm{Tr}_1^n(x)^{p-1}$ is at most p. Expanding $\mathrm{Tr}_1^n(ux)\mathrm{Tr}_1^n(x)^{p-1}$ by the definition of trace function, we can verify that there is a term $(u^p - u)x^{p+(p-1)}$ in $f(x)$. Note that $u^p - u \neq 0$ beause $u \in \mathbb{F}_{p^n} \setminus \mathbb{F}_p$. This implies that the algebraic degree of the p-ary bent function defined by (4.3) is equal to p. Up to the authors' knowledge, this is the first class of p-ary bent functions of algebraic degree p. In other words, we proved that for any odd prime number p, there is a bent function of algebraic degree p. For an odd prime p, some known classes of p-ary monomial and binomial bent functions were listed in Table II [18] and Table 1 [34] respectively. Note that algebraic degree is an affine invariant. According to Table II [18] and Table 1 [34], we conclude that there exist bent functions of the form as in (4.3) which are affinely inequivalent to all known ones listed in Table II [18] and Table 1 [34].

Example 4. *Let $p = 7$, $k = 2$ and $n = 4$. Let α be the generator of $\mathbb{F}_{5^6}^*$ with $\alpha^4 + 5\alpha^2 + 4\alpha + 3 = 0$. Take $\lambda = \alpha^{200}$, $u = \alpha^{90}$. Then f defined by (4.3) is*

$$f(x) = \mathrm{Tr}_1^2(\alpha^{200}x^{50}) - \mathrm{Tr}_1^4(\alpha^{90}x)\mathrm{Tr}_1^4(x)^6.$$

Using Magma, we can verify that $\mathrm{Tr}_1^2(\lambda^{-1}) = 0$, $\mathrm{Tr}_1^4(\lambda^{-1}u) = 0$ and f is bent, which is consistent with the results given in Theorem 4.

Similar to above, by Lemmas 3 and 6, we can obtain another calss of p-ary bent functions of algebraic degree p.

Theorem 5. *For any positive integer $n > 2$, let $\lambda \in \mathbb{F}_{p^n}^*$ and $u \in \mathbb{F}_{p^n} \setminus \mathbb{F}_p$. Then the p-ary function f defined as*

$$f(x) = \mathrm{Tr}_1^n(\lambda x^2) - \mathrm{Tr}_1^n(ux)\mathrm{Tr}_1^n(x)^{p-1} \tag{4.5}$$

is a weakly regular bent function if $\mathrm{Tr}_1^n\left(\frac{1}{4\lambda}\right) = 0$ and $\mathrm{Tr}_1^n\left(\frac{u}{4\lambda}\right) = 0$.

Remark 4. *When $n = 2$, let α be the generator of $\mathbb{F}_{p^2}^*$. The equation $\mathrm{Tr}_1^2\left(\frac{1}{4x}\right) = \frac{1}{4x} + (\frac{1}{4x})^p = 0$, i.e., $x^{p-1} = -1$ has $p - 1$ solutions $\alpha^{\frac{(p+1)i}{2}}$ in $\mathbb{F}_{p^2}^*$, where $i = 2t - 1$ for $1 \leq t \leq p - 1$. For any solution λ of $x^{p-1} = -1$, the equation $\mathrm{Tr}_1^n\left(\frac{y}{4\lambda}\right) = 0$, i.e., $\frac{y}{4\lambda} + (\frac{y}{4\lambda})^p = \frac{y}{4\lambda} + \frac{1}{4\lambda}(\frac{1}{\lambda})^{p-1}y^p = \frac{y}{4\lambda} - \frac{1}{4\lambda}y^p = 0$ leads to $y \in \mathbb{F}_p$, which is contradict with our assumption $u \in \mathbb{F}_{p^n} \setminus \mathbb{F}_p$. Therefore we assume that $n > 2$ in Theorem 5.*

Example 5. *Let $p = 5$ and $n = 3$. Let α be the generator of $\mathbb{F}_{5^3}^*$ with $\alpha^3 + 3\alpha + 3 = 0$. Take $\lambda = \alpha^9$, $u = \alpha^{14}$. Then f defined by (4.5) is*

$$f(x) = \mathrm{Tr}_1^3(\alpha^9x^2) - \mathrm{Tr}_1^3(\alpha^{14}x)\mathrm{Tr}_1^3(x)^4.$$

Using Magma, we can verify that $\mathrm{Tr}_1^3((4\lambda)^{-1}) = 0$, $\mathrm{Tr}_1^3((4\lambda)^{-1}u) = 0$ and f is bent, which is consistent with the results given in Theorem 5.

5. CONCLUSION

The main purpose of this paper is to provide constructions of p-ary functions with low Walsh spectra. We succeed in constructing more quadratic ternary bent, near-bent and 2-plateaued functions, and determining the distribution of the Walsh spectrum of some 2-plateaued functions constructed in this paper. In addition, two classes of non-quadratic p-ary bent functions are obtained. Furthermore, our computer experiments show that there are more p-ary bent functions of the form (1.3) which may be obtained via p-ary Kasami and Sidelnikov monomials, such as $f_1(x) = \mathrm{Tr}_1^4(\xi^6 x^2) + 2\mathrm{Tr}_1^4(x)^2$, $f_2(x) = \mathrm{Tr}_1^4(\xi x^{26}) + 2\mathrm{Tr}_1^4(x)^2$ over $\mathbb{F}_{5^4}$, where ξ is a primitive element of $\mathbb{F}_{5^4}$. Unfortunately, the bentness of theses functions can not be characterized by our method. We leave this for future work.

Acknowledgement

The work of this paper was supported by the National Natural Science Foundation of China (Grants 11371011 and 61403157), the Natural Science Foundation of the Anhui Higher Education Institutions of China (Grants KJ2013B256 and KJ2013B261) and the Natural Science Foundation of the Huainan Normal University (Grant 2013XJ67).

6. Appendix

Computation of the value distribution of the Walsh transform of 2-plateaued function f given by (3.2).

The condition that $\mathrm{Tr}_1^k(\lambda^{-1}u^{3^k+1}) = \mathrm{Tr}_1^k(\lambda^{-1}v^{3^k+1}) = 0$ and $\mathrm{Tr}_1^n(\lambda^{-1}u^{3^k}v) = 2$ leads to $u \neq \pm v$. Otherwise, assume that $u = \pm v$, then $\mathrm{Tr}_1^n(\lambda^{-1}u^{3^k}v) = \pm\mathrm{Tr}_1^n(\lambda^{-1}u^{3^k+1}) = \pm\mathrm{Tr}_1^k(\mathrm{Tr}_k^n(\lambda^{-1}u^{3^k+1})) = \pm 2\mathrm{Tr}_1^k(\lambda^{-1}u^{3^k+1}) = 2$, and thus, $\mathrm{Tr}_1^k(\lambda^{-1}u^{3^k+1}) = \pm 1$. This is contrary to the assumption that $\mathrm{Tr}_1^k(\lambda^{-1}u^{3^k+1}) = 0$.

Let $\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1}) = c_0$ and denote by N_i the number of $a \in \mathbb{F}_{3^n}$ such that $\widehat{\chi}_f(a) = -3^{k+1}\omega^i$, where $i = 0, 1, 2$. The equality (3.7) implies that if $\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1}) = 0$, $\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u) = 0$ and $\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v) = 0$, then $\widehat{\chi}_f(a) = -3^{k+1}$. Hence we have

$$\begin{aligned}
N_0 =& \frac{1}{27}\sum_{a\in\mathbb{F}_{3^n}}\sum_{x\in\mathbb{F}_3}\omega^{x\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})}\sum_{y\in\mathbb{F}_3}\omega^{y\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)}\sum_{z\in\mathbb{F}_3}\omega^{z\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)}\\
=& \frac{1}{27}\sum_{a\in\mathbb{F}_{3^n}}(1+\omega^{\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})}+\omega^{-\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})})(1+\omega^{\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)}\\
&+\omega^{-\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)})(1+\omega^{\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)}+\omega^{-\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)})\\
=& \frac{1}{27}\sum_{a\in\mathbb{F}_{3^n}}(1+\omega^{\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})}+\omega^{-\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})})\big(1+\omega^{\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)}\\
&+\omega^{-\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)}+\omega^{\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)}+\omega^{\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)+\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)}\\
&+\omega^{\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)-\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)}+\omega^{-\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)}+\omega^{-\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)+\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)}\\
&+\omega^{-\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)-\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)}\big).
\end{aligned} \tag{6.1}$$

Recalling $u, v \in \mathbb{F}_{3^n}^*$ and $u \neq \pm v$, we have

$$\begin{aligned}
\sum_{a\in\mathbb{F}_{3^n}}\omega^{\pm\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)} &= \sum_{a\in\mathbb{F}_{3^n}}\omega^{\pm\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)} = \sum_{a\in\mathbb{F}_{3^n}}\omega^{-\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}(u\pm v))}\\
&= \sum_{a\in\mathbb{F}_{3^n}}\omega^{-\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}(v\pm u))} = 0.
\end{aligned} \tag{6.2}$$

From (6.2), the equality (6.1) can be simplified to

$$\begin{aligned}N_0 =& 3^{n-3} + \frac{1}{27}\sum_{a\in\mathbb{F}_{3^n}} (\omega^{\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})} + \omega^{-\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})})\big(1 + \omega^{\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)} \\ &+ \omega^{-\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)} + \omega^{\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)} + \omega^{\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)+\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)} \\ &+ \omega^{\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)-\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)} + \omega^{-\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)} + \omega^{-\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)+\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)} \\ &+ \omega^{-\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)-\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)}\big).\end{aligned}\tag{6.3}$$

Since raising elements of $\mathbb{F}_{3^n}^*$ to the power of 3^k+1 is a 3^k+1 to 1 mapping on to $\mathbb{F}_{3^k}^*$,

$$\begin{aligned}\sum_{a\in\mathbb{F}_{3^n}} \omega^{\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})} &= \sum_{a\in\mathbb{F}_{3^n}} \omega^{-\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})} \\ &= 1 + (3^k+1)\sum_{b\in\mathbb{F}_{3^k}^*} \omega^{\mathrm{Tr}_1^k(\lambda^{-1}b)} = -3^k\end{aligned}\tag{6.4}$$

because $\lambda \in \mathbb{F}_{3^k}^*$.

From $\mathrm{Tr}_1^k(\lambda^{-1}u^{3^k+1}) = 0$, we obtain

$$\begin{aligned}&\sum_{a\in\mathbb{F}_{3^n}} \omega^{\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})\pm\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)} \\ =& \sum_{a\in\mathbb{F}_{3^n}} \omega^{\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})\pm\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)+\mathrm{Tr}_1^k(\lambda^{-1}u^{3^k+1})} \\ =& \sum_{a\in\mathbb{F}_{3^n}} \omega^{\mathrm{Tr}_1^k(\lambda^{-1}(a\pm u)^{3^k+1})} = -3^k\end{aligned}\tag{6.5}$$

and

$$\sum_{a\in\mathbb{F}_{3^n}} \omega^{-\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})\pm\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)} = -3^k.\tag{6.6}$$

Similarly, it follows from $\mathrm{Tr}_1^k(\lambda^{-1}v^{3^k+1}) = 0$ that

$$\begin{aligned}&\sum_{a\in\mathbb{F}_{3^n}} \omega^{\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})\pm\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)} \\ =& \sum_{a\in\mathbb{F}_{3^n}} \omega^{\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})\pm\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)+\mathrm{Tr}_1^k(\lambda^{-1}v^{3^k+1})} \\ =& \sum_{a\in\mathbb{F}_{3^n}} \omega^{\mathrm{Tr}_1^k(\lambda^{-1}(a\pm v)^{3^k+1})} = -3^k\end{aligned}\tag{6.7}$$

and

$$\sum_{a\in\mathbb{F}_{3^n}} \omega^{-\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})\pm\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)} = -3^k.\tag{6.8}$$

On the other hand, it follows from $\mathrm{Tr}_1^k(\lambda^{-1}u^{3^k+1})=0$, $\mathrm{Tr}_1^k(\lambda^{-1}v^{3^k+1})=0$ and $\mathrm{Tr}_1^n(\lambda^{-1}u^{3^k}v)=2$ that

$$\begin{aligned}
&\sum_{a\in\mathbb{F}_{3^n}}\omega^{\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})+\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)+\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)}\\
=&\omega\sum_{a\in\mathbb{F}_{3^n}}\omega^{\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})+\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)+\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)+\mathrm{Tr}_1^n(\lambda^{-1}u^{3^k}v)}\\
&\times\omega^{\mathrm{Tr}_1^k(\lambda^{-1}u^{3^k+1})+\mathrm{Tr}_1^k(\lambda^{-1}v^{3^k+1})}\\
=&\omega\sum_{a\in\mathbb{F}_{3^n}}\omega^{\mathrm{Tr}_1^k(\lambda^{-1}(a+v+u)^{3^k+1})}\\
=&-3^k\omega.
\end{aligned}\tag{6.9}$$

Similarly, we have

$$\begin{aligned}
\sum_{a\in\mathbb{F}_{3^n}}\omega^{\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})+\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)-\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)}=&\omega^2\sum_{a\in\mathbb{F}_{3^n}}\omega^{\mathrm{Tr}_1^k(\lambda^{-1}(a-v+u)^{3^k+1})}\\
=&-3^k\omega^2,
\end{aligned}\tag{6.10}$$

$$\begin{aligned}
\sum_{a\in\mathbb{F}_{3^n}}\omega^{\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})-\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)+\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)}=&\omega^2\sum_{a\in\mathbb{F}_{3^n}}\omega^{\mathrm{Tr}_1^k(\lambda^{-1}(a+v-u)^{3^k+1})}\\
=&-3^k\omega^2
\end{aligned}\tag{6.11}$$

and

$$\begin{aligned}
\sum_{a\in\mathbb{F}_{3^n}}\omega^{\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})-\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)-\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)}=&\omega\sum_{a\in\mathbb{F}_{3^n}}\omega^{\mathrm{Tr}_1^k(\lambda^{-1}(a-v-u)^{3^k+1})}\\
=&-3^k\omega.
\end{aligned}\tag{6.12}$$

Using $\mathrm{Tr}_1^k(\lambda^{-1}u^{3^k+1})=0$, $\mathrm{Tr}_1^k(\lambda^{-1}v^{3^k+1})=0$ and $\mathrm{Tr}_1^n(\lambda^{-1}u^{3^k}v)=2$ again, we obtain

$$\begin{aligned}
&\sum_{a\in\mathbb{F}_{3^n}}\omega^{-\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})+\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)+\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)}\\
=&\omega^2\sum_{a\in\mathbb{F}_{3^n}}\omega^{-\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})+\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)+\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)-\mathrm{Tr}_1^n(\lambda^{-1}u^{3^k}v)}\\
&\times\omega^{-\mathrm{Tr}_1^k(\lambda^{-1}u^{3^k+1})-\mathrm{Tr}_1^k(\lambda^{-1}v^{3^k+1})}\\
=&\omega^2\sum_{a\in\mathbb{F}_{3^n}}\omega^{\mathrm{Tr}_1^k(-\lambda^{-1}(a-v-u)^{3^k+1})}\\
=&-3^k\omega^2.
\end{aligned}\tag{6.13}$$

Similarly, we have

$$\begin{aligned}
\sum_{a\in\mathbb{F}_{3^n}}\omega^{-\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})+\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)-\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)}=&\omega\sum_{a\in\mathbb{F}_{3^n}}\omega^{\mathrm{Tr}_1^k(-\lambda^{-1}(a+v-u)^{3^k+1})}\\
=&-3^k\omega,
\end{aligned}\tag{6.14}$$

$$\begin{aligned}\sum_{a\in\mathbb{F}_{3^n}} \omega^{-\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})-\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)+\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)} =&\omega \sum_{a\in\mathbb{F}_{3^n}} \omega^{\mathrm{Tr}_1^k(-\lambda^{-1}(a-v+u)^{3^k+1})}\\ =&-3^k\omega \end{aligned}\tag{6.15}$$

and

$$\begin{aligned}\sum_{a\in\mathbb{F}_{3^n}} \omega^{\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})-\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)-\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)} =&\omega^2 \sum_{a\in\mathbb{F}_{3^n}} \omega^{\mathrm{Tr}_1^k(\lambda^{-1}(a-v-u)^{3^k+1})}\\ =&-3^k\omega^2. \end{aligned}\tag{6.16}$$

Combining (6.3)-(6.16), we get

$$N_0 = 3^{n-3} + \frac{1}{27}(-10\cdot 3^k - 4\cdot 3^k(\omega+\omega^2)) = 3^{n-3} - \frac{1}{27}6\cdot 3^k = 3^{n-3} - 2\cdot 3^{k-2}.$$

We next compute N_1 and N_2. According to (3.7), if $\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1}) = 2$, $\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u) = 0$ and $\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v) = 0$, then $\widehat{\chi}_f(a) = -3^{k+1}\omega$. Therefore we have

$$\begin{aligned}N_1 =&\frac{1}{27}\sum_{a\in\mathbb{F}_{3^n}}\sum_{x\in\mathbb{F}_3}\omega^{x(\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})-2)}\sum_{y\in\mathbb{F}_3}\omega^{y\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)}\sum_{z\in\mathbb{F}_3}\omega^{z\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)}\\
=&\frac{1}{27}\sum_{a\in\mathbb{F}_{3^n}}(1+\omega^{1+\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})}+\omega^{2-\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})})(1+\omega^{\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)}\\
&+\omega^{-\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)})(1+\omega^{\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)}+\omega^{-\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)})\\
=&\frac{1}{27}\sum_{a\in\mathbb{F}_{3^n}}(1+\omega^{1+\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})}+\omega^{2-\mathrm{Tr}_1^k(\lambda^{-1}a^{3^k+1})})\big(1+\omega^{\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)}\\
&+\omega^{-\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)}+\omega^{\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)}+\omega^{\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)+\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)}\\
&+\omega^{\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)-\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)}+\omega^{-\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)}+\omega^{-\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)+\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)}\\
&+\omega^{-\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}u)-\mathrm{Tr}_1^n(\lambda^{-1}a^{3^k}v)}\big).\end{aligned}$$

By a similar process of calculating N_0, we can obtain

$$\begin{aligned}N_1 =&3^{n-3}+\frac{1}{27}(-5\cdot 3^k\omega - 5\cdot 3^k\omega^2 - 2\cdot 3^k\omega(\omega+\omega^2) - 2\cdot 3^k\omega^2(\omega+\omega^2))\\
=&3^{n-3}+\frac{1}{27}(-3\cdot 3^k(\omega+\omega^2)) = 3^{n-3}+3^{k-2}.\end{aligned}$$

According to Parseval equation $\sum_{a\in\mathbb{F}_{3^n}}|\widehat{\chi}_f(a)|^2 = 3^{2n}$, we get the number of $a\in\mathbb{F}_{3^n}$ such that $|\widehat{\chi}_f(a)| = 3^{k+1}$ is equal to 3^{n-2} and the number of $a\in\mathbb{F}_{3^n}$ such that $\widehat{\chi}_f(a) = 0$ is equal to $3^n - 3^{n-2}$. This implies that $N_2 = 3^{n-3}+3^{k-2}$.

REFERENCES

[1] A. Canteaut, P. Charpin, G. M. Kyureghyan, A new class of monomial bent functions, Finite Fields Appl. **14**(1) (2008) 221–241.

[2] C. Carlet, Two new classes of bent functions, In: Advances in cryptology-EUROCRYPT93, Springer, 1994, pp. 77–101.

[3] C. Carlet, On bent and highly nonlinear balanced/resilient functions and their algebraic immunities. In: Applied Algebra, Algebraic Algorithms and Error-Correcting Codes, Springer, 2006, pp. 1–28.

[4] C. Carlet, Boolean functions for cryptography and error correcting codes, in Boolean Models and Methods in Mathematics, Computer Science, and Engineering, Y. Crama and P. L. Hammer, Eds. Cambridge, U.K.: Cambridge Univ. Press, 2010, pp. 257-397.

[5] C. Carlet, E. Prouff, On plateaued functions and their constructions, in Proceedings of Fast Software Encryption (FSE), Lecture Notes in Computer Science, vol. 2887 (2003), pp. 54–73.

[6] C. Carlet, S. Mesnager, On Dillon's class $\mathcal{H}$ of bent functions, Niho bent functions and o-polynomials. J. Combin. Theory Ser A **118**(8) (2011) 2392–2410.

[7] A. Çeşmelioğlu, W. Meidl, A. Pott, Generalized Maiorana-McFarland class and normality of p-ary bent functions, Finite Fields Appl. **24**(9) (2013) 105-117.

[8] A. Çeşmelioğlu, G. McGuire, W. Meidl, A construction of weakly and non-weakly regular bent functions, J. Combin. Theory Ser A **119** (2012) 420–429.

[9] P. Charpin, E. Pasalic, C. Tavernier, On bent and semi-bent quadratic Boolean functions, IEEE Trans. Inf. Theory **51**(12) (2005) 4286–4298.

[10] S. Chee, S. Lee, K. Kim, Semi-bent functions, in Advances in Cryptology-ASIACRYPT'94, Springer, pp. 105-118, 1995.

[11] R. S. Coulter, M. Henderson, The compositional inverse of a class of permutation polynomials over a finite field. Bull. Aust. Math. Soc. **65**(03) (2002) 521–526.

[12] J. F. Dillon, Elementary hadamard difference sets, Ph.D. thesis, University of Maryland, College Park, 1974.

[13] H. Dobbertin, G. Leander, A survey of some recent results on bent functions, In: Sequences and Their Applications-SETA 2004,Springer, 2005, pp. 1–29.

[14] T. Helleseth, Correlation of m-sequences and related topics, In: Sequences and their Applications, Springer, 1999, pp. 49–66.

[15] T. Helleseth, A. Kholosha, Monomial and quadratic bent functions over the finite fields of odd characteristic, IEEE Trans. Inf. Theory **52**(5) (2006) 2018–2032.

[16] T. Helleseth, A. Kholosha, New binomial bent functions over the finite fields of odd characteristic, IEEE Trans. Inf. Theory **56**(9) (2010) 4646–4652.

[17] X. D. Hou, p-Ary and q-ary versions of certain results about bent functions and resilient functions, Finite Fields Appl. **10**(4) (2004) 566–582.

[18] W. Jia, X. Zeng, T. Helleseth, C. Li, A class of binomial bent functions over the finite fields of odd characteristic, IEEE Trans. Inf. Theory **58**(9) (2012) 6054–6063.

[19] K. Khoo, G. Gong, D. R. Stinson, A New Characterization of Semi-bent and Bent Functions on Finite Fields, Des. Codes Cryptogr. **38**(2) (2006) 279–295.

[20] P. V. Kumar, R. A. Scholtz, L. R. Welch, Generalized bent functions and their properties, J. Combin. Theory Ser A **40**(1) (1985) 90–107.

[21] N. Li, T. Helleseth, X. Tang, A. Kholosha, Several new classes of bent functions from Dillon exponents, IEEE Trans. Inf. Theory **59**(3) (2013) 1818–1831.

[22] S. Li, L. Hu, X. Zeng, Constructions of p-ary quadratic Bent functions, Acta Appl. Math. **100**(3) (2008) 227–245.

[23] S. C. Liu, J. J. Komo, Nonbinary Kasami sequences over $\mathbf{GF}(p)$, IEEE Trans. Inf. Theory **38**(4) (1992) 1409–1412.

[24] M. Matsui, Linear cryptanalysis method for DES cipher, In: Advances in Cryptology-EUROCRYPT93, Springer, 1994, pp. 386–397.

[25] S. Mesnager, A new class of bent and hyper-bent Boolean functions in polynomial forms, Des. Codes Cryptogr. **59**(1-3) (2011) 265–279

[26] S. Mesnager, On Semi-bent Functions and Related Plateaued Functions Over the Galois Field $\mathbb{F}_{2^n}$, In: Open Problems in Mathematics and Computational Science, Springer , 2013, pp. 243–273.

[27] S. Mesnager, Several new infinite families of bent functions and their duals, IEEE Trans. Inf. Theory **60**(7) (2014) 4397–4407.

[28] S. Mesnager, Bent functions: fundamentals and results, Springer, 2015, to appear.

[29] O. S. Rothaus,: On bent functions, J. Combin. Theory Ser A **20**(3) (1976) 300–305.

[30] A. Tuxanidy, Q. Wang, Compositional inverses, complete mappings, orthogonal Latin squares and bent functions, arXiv:1409.6540 [math.NT], preprint (2014).

[31] B. Wu, The compositional inverses of linearized permutation binomials over finite fields, arXiv:1311.2154v1 [math.NT], preprint (2013).

[32] B. Wu, Z. Liu, Linearized polynomials over finite fields revisited, Finite Fields Appl **22** (2013) 79–100.

[33] G. Xu, X. Cao, S. Xu, Constructing new APN functions and bent functions over finite fields of odd characteristic via the switching method, Cryptogr. Commun.(2015), DOI:10.1007/s12095-015-0145-6, in press.

[34] D. Zheng, L. Yu, L. Hu, On a class of binomial bent functions over the finite fields of odd characteristic, Appl. Algebra Eng. Commun. Comput. **24**(6) (2013) 461–475.

[35] D. Zheng, L. Yu, L. Hu, Quadratic Bent and Semi-bent Functions over Finite Fields of Odd Characteristic, Chinese J. Electron. **23**(4) (2014) 767–772.

[36] Y. Zheng, X.M. Zhang, Plateaued functions, in Advances in Cryptology ICICS 1999. Lecture Notes in Computer Science, vol. 1726 (Springer, Berlin, 1999), 284–300.

[37] Y. Zheng, X.M. Zhang, Relationships between bent functions and complementary plateaued functions, Lecture Notes in Computer Science, **24** (1999) 60–75.

G. Xu is with the Department of Mathematics, Nanjing University of Aeronautics and Astronautics, Nanjing 210016, China, and also with the Department of Mathematics, Huainan Normal University, Huainan 232038, China (e-mail: `xuguangkuiy@163.com`).

Xiwang Cao is with the School of Mathematical Sciences, Nanjing University of Aeronautics and Astronautics, Nanjing 210016, China, and also with the State Key Laboratory of Information security, Institute of Information Engineering, Chinese Academy of Sciences, Beijing 100093, China (email: `xwcao@nuaa.edu.cn`).